



**UNIVERSITI PUTRA MALAYSIA**

**PARTIAL PURIFICATION AND CHARACTERIZATION OF  
GLUTATHIONE S-TRANSFERASE FROM THE LIVERS OF  
MALAYSIAN CATFISH (CLARIAS BATRACHUS) AND MALAYSIAN  
RED TILAPIA (OREOCHROMIS SP.)**

**CHIN CHEE MENG.**

**FBSB 2005 32**

**PARTIAL PURIFICATION AND CHARACTERIZATION OF GLUTATHIONE  
S-TRANSFERASE FROM THE LIVERS OF MALAYSIAN CATFISH (*CLARIAS  
BATRACHUS*) AND MALAYSIAN RED TILAPIA (*OREOCHROMIS SP.*)**

**By**

**CHIN CHEE MENG**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra  
Malaysia, in Fulfilment of the Requirements for the Degree of  
Master of Science**

**July 2005**



**Dedicated to my parents**

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in  
fulfilment of the requirements for the degree of Master of Science

**PARTIAL PURIFICATION AND CHARACTERIZATION OF GLUTATHIONE S-  
TRANSFERASE FROM THE LIVERS OF MALAYSIAN CATFISH (*CLARIAS*  
*BATRACHUS*) AND MALAYSIAN RED TILAPIA (*OREOCHROMIS SP.*)**

By

**CHIN CHEE MENG**

**July 2005**

**Chairman : Professor Nor Aripin Shamaan, PhD**

**Faculty : Biotechnology and Biomolecular Sciences**

The present study is concerned mainly with partial purification and characterization of cytosolic GST from livers of Malaysian catfish and Malaysian red tilapia as both species are of economic and commercial importance in Malaysia being the major freshwater fishes consumed. This study hopes to establish the patterns of cytosolic glutathione S-transferase isoenzymes in the catfish and red tilapia. This may be useful to further achieve an understanding toward this enzyme in view of using it as a tool in environmental monitoring. The hepatic GST enzyme from catfish and red tilapia was partially purified 15X and 27X respectively in comparison to the ultra-centrifuged cytosolic fraction by affinity chromatography. Specific GST activity of 12.69 unit/mg protein and 33.42 unit/mg protein was obtained from livers of catfish and red tilapia using 1-chloro-2,4-dinitrobenzene (CDNB) as a substrate. Isolation of GST isoenzymes from affinity purified fractions was achieved by preparative isoelectric-focusing. Two isoenzymes; one major isoenzyme designated Ci1 and one minor isoenzyme

designated Ci2 were isolated from catfish liver with an apparent pI of 6.20 and 8.73 respectively. One isoenzyme designated Ti1 was isolated from red tilapia liver with an apparent pI of 9.14. SDS-PAGE analysis suggests that the isolated isoenzymes appear to be homodimeric in nature with subunit molecular weight of  $29.7 \pm 1.7$  kDa (Ci1),  $27.7 \pm 1.3$  kDa (Ci2) and  $29.9 \pm 0.9$  kDa (Ti1). As has been found for most GSTs, highest catalytic activity was obtained with CDNB. With the exception of the isoenzyme from cytosol and affinity purified fractions of red tilapia, none of the catfish fractions displayed enzyme activities towards 1,2-dichloro-4-nitrobenzene (DCNB) and ethacrynic acid (EA). Therefore, both catfish and tilapia possess hepatic glutathione S-transferase activity, indicating that they are capable of conjugating endogenous or xenobiotic metabolites/compounds as a result of foreign exposure or oxidative metabolism with glutathione, thereby making it a useful tool as a effective biomarker of aquatic contamination.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia  
sebagai memenuhi keperluan untuk ijazah Master Sains

**PURIFIKASI SEPARA DAN PENCIRIAN ENZIM GLUTATHIONE S-  
TRANSFERASE DARIPADA HATI IKAN KELI MALAYSIA (*CLARIAS  
BATRACHUS*) DAN IKAN TILAPIA MERAH MALAYSIA (*OREOCHROMIS  
SP.*)**

Oleh

**CHIN CHEE MENG**

**July 2005**

**Pengerusi : Profesor Nor Aripin Shamaan, PhD**

**Fakulti : Bioteknologi dan Sains Biomolekul**

Kajian ini mengenai purifikasi separa dan pencirian GST sitosol daripada ikan keli Malaysia dan ikan tilapia merah Malaysia disebabkan kedua-dua spesies tersebut mempunyai kepentingan ekonomi dan komersial di Malaysia memandangkan ia adalah ikan air tawar yang sering dimakan. Kajian ini diharapkan dapat menentukan corak isoenzim sitosol GST dalam hati ikan keli dan tilapia merah. Ini mungkin berguna untuk mencapai pemahaman ke atas enzim ini dimana ia boleh dijadikan sebagai alat untuk memantau persekitaran. Enzim GST hepatik daripada ikan keli dan tilapia merah telah disepara-tulen sebanyak 15X dan 27X masing-masing dibandingkan dengan fraksi sitosol ultra-centrifuged oleh kromatografi afiniti. Aktiviti spesifik enzim yang diperolehi adalah 12.69 unit/mg dan 33.42 unit/mg daripada hati ikan keli dan tilapia merah masing-masing dengan menggunakan 1-chloro-2,4-dinitrobenzene (CDNB) sebagai substrat. Pemencilan isoenzim GST daripada fraksi yang ditulenkan dengan kromatografi afiniti dicapai dengan menggunakan "preparative

isoelectric-focusing". Dua isoenzim; satu isoenzim major Ci1 dan satu isoenzim minor Ci2 dipencilkan daripada hati ikan keli dengan nilai pI 6.20 dan 8.73 masing-masing. Satu isoenzim Ti1 dipencilkan daripada hati ikan tilapia merah dengan nilai pI 9.14. Analisis SDS-PAGE mencadangkan isoenzim-isoenzim yang dipencilkan mempunyai struktur homodimer dengan berat molekul subunit  $29.7 \pm 1.7$  kDa (Ci1),  $27.7 \pm 1.3$  kDa (Ci2) dan  $29.9 \pm 0.9$  kDa (Ti1). Seperti yang ditemui dengan kebanyakan GST yang lain, aktiviti katalitik yang tertinggi diperolehi dengan CDNB. Dengan pengecualian isoenzim daripada sitosol dan fraksi yang ditulen oleh kromatografi afiniti daripada ikan tilapia merah, tiada fraksi ikan keli yang menunjukkan aktiviti dengan 1,2-dichloro-4-nitrobenzene (DCNB) dan ethacrynic acid (EA). Oleh itu, kedua-dua ikan keli dan tilapia mempunyai aktiviti glutathione S-transferase, menunjukkan bahawa mereka mampu mengkonjugasikan metabolik/bahan endogenus atau xenobiotik akibat daripada terdedah kepada persekitaran atau metabolisme oksidatif dengan glutathione, oleh itu membuatnya alat yang berguna sebagai petanda biologi untuk pencemaran akuatik.

## ACKNOWLEDGEMENTS

First and foremost I would like to express my heartfelt gratitude and appreciation to my supervisor, Prof. Nor Aripin Shamaan for his advice, guidance, ideas and critical discussions throughout the course of this project. He was always available for consultation even after office hours and has never failed to give an encouraging word when in need. Special thanks are extended to the other members of the Supervisory Committee, Dr. Yunus and Encik Ismail Omar for their ideas, advice and guidance given whenever sought.

I would also like to thank Encik Ibrahim and Encik Shamsudin for their help in preparing the chemicals and equipments needed during the course of the project. Not forgetting also to my lab members; Eddie Chia, Thong Chuan, Che Wan, Lan, Kak Neni, and Nina for their cooperation, companionship in the lab till the wee hours of morning and their help in supplying chemicals and lab equipments. A million thanks to them!

I wish to thank the Ministry of Science and Technology for the IRPA grant that funded this project and also for the PASCA scholarship given to me for two years.

A big thank you is also expressed to my parents for their encouragement and financial support to me throughout my study.



I certify that an Examination Committee met on 20<sup>th</sup> July 2005 to conduct the final examination of Chin Chee Meng on his Master of Science thesis entitled “Partial Purification and Characterization of Glutathione S-Transferase from the Liver of Malaysian Catfish (*Clarias batrachus*) and Malaysian Red Tilapia (*Oreochromis* sp.)” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

**Juzu Hayati Arshad, PhD**

Associate Professor

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Chairman)

**Mohd Arif Syed, PhD**

Professor

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Internal Examiner)

**Muhajir Hamid, PhD**

Lecturer

Faculty of Biotechnology and Biomolecular Sciences

Universiti Putra Malaysia

(Internal Examiner)

**Yasmin Anum Mohd Yusof, PhD**

Associate Professor

Faculty of Medicine

Universiti Kebangsaan Malaysia

(External Examiner)



**GULAM RUSUL RAHMAT ALI, PhD**

Professor/Deputy Dean

School of Graduate Studies

Universiti Putra Malaysia

Date:

**22 AUG 2005**

This thesis submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfilment of the requirements for the degree of Master of Science. The members of the Supervisory Committee are as follows :

**Nor Aripin Shamaan, PhD**


Professor  
Faculty of Biotechnology and Biomolecular Sciences  
Universiti Putra Malaysia  
(Chairman)

**Mohd Yunus Abdul Shukur, PhD**

Lecturer  
Faculty of Biotechnology and Biomolecular Sciences  
Universiti Putra Malaysia  
(Member)

**Ismail Omar, M.Sc.**

Lecturer  
Faculty of Biotechnology and Biomolecular Sciences  
Universiti Putra Malaysia  
(Member)



---

**AINI IDERIS, PhD**  
Professor/Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date : 08 SEP 2005

**DECLARATION**

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.

  
CHIN CHEE MENG

Date : 12/8/2005

## TABLE OF CONTENTS

	<b>Page</b>
<b>DEDICATION</b>	ii
<b>ABSTRACT</b>	iii
<b>ABSTRAK</b>	v
<b>ACKNOWLEDGEMENT</b>	vii
<b>APPROVAL</b>	viii
<b>DECLARATION</b>	x
<b>LIST OF TABLES</b>	xiii
<b>LIST OF FIGURES</b>	xiv
<b>LIST OF ABBREVIATIONS</b>	xvi
 <b>CHAPTER</b>	
 <b>I</b>	
<b>INTRODUCTION</b>	19
Objectives	22
 <b>II</b>	
<b>LITERATURE REVIEW</b>	23
The Enzymes of Detoxification	23
Glutathione	24
Mercapturic acid	26
Glutathione S-Transferase	26
The Discovery of Glutathione S-Transferase	33
Isoenzymes of Glutathione S-Transferase	34
Cellular location of Glutathione S-Transferase	36
Nomenclature	37
The Final Classification	42
The Substrates of Glutathione S-Transferase	43
Assay Methods	48
Spectrophotometric Method	48
Titrimetric Method	49
Colorimetric Method	49
Radioisotope Method	51
Thiolysis Method	51
Purification Studies	51
Properties of Glutathione S-Transferase in Fish	55
The Structure of Glutathione S-Transferase	60
Kinetic Mechanism of Glutathione S-Transferase	66
Induction of Glutathione S-Transferase Activity	68
Inhibition of Glutathione S-Transferase Activity	71
Glutathione S-Transferase as Biomarkers	72
The Multifunctional Nature of Glutathione S-Transferase	73
Evolution of Glutathione S-Transferase	79



<b>III</b>	<b>METHODOLOGY</b>	<b>81</b>
	Tissue Samples	81
	Determination of the Enzymatic Activity	81
	Measurement of Protein Concentration	83
	Preparation of the Cytosol	83
	Cytosol Purification by Glutathione-Agarose Affinity Chromatography	84
	Isoelectric-focusing	85
	Denaturing SDS-Polyacrylamide Gel Electrophoresis	88
	Determination of Substrate Specificity	89
<b>IV</b>	<b>RESULTS</b>	<b>90</b>
	Preparation of Cytosol	90
	Purification of Cytosol by Affinity Chromatography	91
	Effects of High Concentration of Salt	95
	Isoelectric-focusing	98
	Repeat of the Purification Run	104
	SDS-PAGE Analysis	109
	Substrate Specificity	115
	Summary of Purification and Characterization of Glutathione S-Transferase From the Livers of Catfish and Red Tilapia	117
<b>V</b>	<b>DISCUSSION</b>	<b>119</b>
<b>VI</b>	<b>CONCLUSION</b>	<b>132</b>
	<b>BIBLIOGRAPHY</b>	<b>134</b>
	<b>APPENDIX</b>	<b>159</b>
	<b>BIODATA OF THE AUTHOR</b>	<b>160</b>

## LIST OF TABLES

Table		Page
1	Human members of the MAPEG family	29
2	Classification of rat glutathione S-transferases	41
3	Conditions for spectrophotometric assays	50
4	Quaternary structure of rat GST	63
5	Formation of the light and dense pH gradient	87
6	Preparation of catfish cytosol	90
7	Preparation of red tilapia cytosol	90
8	Purification of catfish cytosol by affinity chromatography	91
9	Purification of red tilapia cytosol by affinity chromatography	91
10	Effect of high salt concentration on the purification procedure	95
11	Purification of catfish cytosol by affinity chromatography (repeat experiment)	104
12	Purification of red tilapia cytosol by affinity chromatography (repeat experiment)	104
13	GST activities of isolated isoenzymes and isolated electro-focused peaks from affinity purified samples and cytosol fractions from catfish and red Tilapia toward different substrates	116
14	Summary of purification procedures and quantitative data of hepatic glutathione S-transferases from The Malaysian Catfish ( <i>Clarius batrachus</i> ) and Malaysian Red Tilapia ( <i>Oreochromis sp.</i> )	118

## LIST OF FIGURES

Figure		Page
1	The mercapturic acid pathway for urinary excretion of glutathione conjugate	27
2	Affinity-chromatographic purification of catfish livers on glutathione-agarose matrix	92
3	Affinity-chromatographic purification of red tilapia livers on glutathione-agarose matrix	93
4	Affinity-chromatographic purification of catfish livers on glutathione-agarose matrix using 1M NaCl eluting buffer	97
5	Isoelectric-focusing profile of affinity purified samples from catfish liver	100
6	Isoelectric-focusing profile of affinity purified samples from red tilapia liver	101
7	Isoelectric-focusing profile of cytosol samples from catfish liver	102
8	Isoelectric-focusing profile of cytosol samples from red tilapia liver	103
9	Affinity-chromatographic purification (repeat experiment) of catfish liver on glutathione-agarose matrix	106
10	Affinity-chromatographic purification (repeat experiment) of red tilapia liver on glutathione-agarose matrix	107
11	Isoelectric-focusing profile of affinity purified samples (repeat experiment) from catfish liver	108
12	SDS-PAGE analysis of electrofocused samples from affinity purified pooled fractions of catfish and red tilapia livers	110
13	SDS-PAGE analysis of electrofocused samples from cytosolic fractions of catfish liver	111

14	SDS-PAGE analysis of electrofocused samples from cytosolic fractions of red tilapia liver	112
15	SDS-PAGE analysis of electrofocused samples from affinity purified pooled fractions (first run and repeat experiment) of catfish and (repeat experiment) of red tilapia livers	114



## LIST OF ABBREVIATIONS

$\alpha$	alpha
$\beta$	beta
Br	bromine
BSA	bovine serum albumin
CDNB	1-chloro-2,4-dinitrobenzene
Cl	chlorine
C-terminal	carboxyl-terminal
Cys	cysteine
$\Delta\epsilon$	molar extinction coefficient
DCNB	1,2-dichloro-4-nitrobenzene
DDT	dichlorodiphenyltrichloroethane
DNA	deoxyribonucleic acid
$\epsilon$	epsilon
E.C.	Enzyme Commission
EA	ethacrynic acid
EDTA	ethylenediaminetetraacetic acid
EST	expressed sequence tag
GDP	gross domestic product
Glu	glutamate
Gly	glycine
GS <sup>-</sup>	glutathione thiolate ion
GSH	reduced glutathione

G-site	glutathione binding site
GST	glutathione S-transferase
HPLC	high performance liquid chromatography
H-site	hydrophobic binding site
I	iodine
KCl	potassium chloride
kDa	kilo Dalton
M	molar
MAPEG	Membrane Associated Proteins in Eicosanoid and Glutathione metabolism
mol	mole
MPOB	Malaysian Palm Oil Board
mRNA	messenger RNA
MW	molecular weight
NaCl	sodium chloride
N-terminal	amino-terminal
$\theta$	theta
$\rho$	rho
PhIP	2-amino-1-methyl-6-phenylimidazo[4,5- <i>b</i> ]pyridine
RNA	ribonucleic acid
SDS-PAGE	sodium dodecyl sulfate polyacrylamide gel electrophoresis
-SH	sulfurhydryl group
$\pi$	pi
TEMED	N, N, N', N'-tetramethylethylenediamine

$\mu$	mu
U	unit
UPM	Universiti Putra Malaysia
$\gamma$	gamma

## CHAPTER I

### INTRODUCTION

The Malaysian fisheries sector plays an important role in providing food and as a source of protein to the general public. In 2000, it contributed about 1.60% to the national Gross Domestic Product (GDP)<sup>a</sup> and provided direct employment to 82,000 fishermen and 22,000 fish culturists. With the recent episodes of Nipah virus infecting pigs and the bird flu attack in chickens, more people are turning to other sources of proteins such as fish. Therefore, the Department of Fisheries is taking several measures in order to provide a sustained production of fish in the country. Although deep sea fishing still remains as the easiest method for fish production, however water pollution and over-fishing are taking its toll on the fish population in the wild causing shortages and resulting in the increasing prices of fish. Therefore, the real potential lies in the aquaculture industry in order to achieve a sustainable growth in fish production.

Freshwater aquaculture contributes a significant percentage (30.2%) to the total aquaculture production in 2000. Some of the major freshwater species cultured were freshwater Malaysian Catfish (*Clarius batrachus*) and Red Tilapia (*Oreochromis sp.*).

---

<sup>a</sup> National GDP RM 339.42 billion (Statistics Handbook, Statistics Department Malaysia, 2001)

The catfish are demersal creatures with the ability to survive in muddy and polluted water. Generally, they are found in rivers and swampy areas. They are omnivorous in nature surviving on water insects, worms, shrimps, snails and rotting materials such as plants. They have a special breathing system which allows them to survive for short periods of time on land. In Malaysia the Malaysian catfish is usually mated with the African catfish (*Clarias gariepinus*) as the African catfish displays higher growth rate and can attain a size of 1 kg in 6 months. The red tilapia can be found in freshwater, saltwater and also brackish water bodies. They are also omnivorous, surviving on a diet of worms, plants and small fish (Basir and Daud, 1994).

As the two species represent an economic importance in Malaysia, maximum yield during the breeding and production are of utmost importance to maximize profits. Therefore, there is a need for a monitoring system to monitor the state of the fish habitat and the stress levels of the fish (i.e overcrowding, parasite infections).

Glutathione S-Transferases (GSTs, E.C. 2.5.1.18) are a multigene family of multifunctional enzymes which can be found in bacteria, yeasts, higher plants, mollusks, crustaceans, insects, fish, amphibians and mammals (Jakoby, 1978; Mannervik, 1985b). They catalyze the nucleophilic attack of the sulfur atom of glutathione on compounds possessing an electrophilic center (Hayes and Pulford, 1995). Therefore, GSTs play a key role in the cellular detoxification process in by conjugating a broad spectrum of endobiotic and xenobiotic electrophilic substrates (as a result of

environmental exposure or oxidative metabolism) with intracellular glutathione into more readily excretable products (Mannervik and Danielson, 1988; Hayes and Pulford, 1995; Armstrong, 1997).

Certain GST isoenzymes display selective response towards pollutants and high activity towards certain substrates, making it a useful tool as an effective biomarker of aquatic contamination (Egaas et al., 1999; Gallagher et al., 1996).

## Objectives

The objectives of the present study are to:-

1. Partially purified the cytosolic glutathione S-transferase (E.C. 2.5.1.18) from the livers of Malaysian catfish (*Clarius batrachus*) and red tilapia (*Oreochromis sp.*)
2. Partially characterized the cytosolic glutathione S-transferase from the livers of Malaysian catfish (*Clarius batrachus*) and red tilapia (*Oreochromis sp.*)

Ultimately, this study hopes to establish the patterns of the cytosolic glutathione S-transferases (i.e. the dimeric structures, specific activities, pI values, molecular weight and substrate specificity) in the catfish and red tilapia. This may be useful to further achieve an understanding toward this enzyme in view of using it as a tool in environmental monitoring.

## CHAPTER II

### LITERATURE REVIEW

#### The Enzymes of Detoxification

All living organism are continuously subjected to foreign external (xenobiotics) compounds; naturally occurring or man-made such as chemicals in the air, water, food additions and drugs or endogenous toxicants produced as a result of the metabolism of hormones, steroids and oxidative metabolism. Most of these compounds are lipophilic and might be excreted slowly from the body. Therefore, the toxicity of the compounds is increased as a result of the accumulation in the tissues of the organism. Given these exposures, the ability to detoxify is important to the overall health and the survivability of the organism.

The process of detoxification through which the non-polar (lipophilic) toxins are converted to polar (hydrophilic) compounds that can be readily excreted by the organism occurs in two classical steps known as Phase 1 and Phase 2. Each of the phases involved the so-called detoxification enzymes that have broad specificity towards a wide variety of toxins faced by the organism. The two phases are interconnected; Phase 1 reactions normally generate intermediate end-products that are toxic to the tissues. Phase 2 reactions conjugates the end-products of Phase 1 reactions to yield final products that are eliminated from the organism. Phase 1 reactions are characterized by the cytochrome P450 supergene family of isoenzymes that catalyzed the addition



of a reactive group to the substrate (toxins). However the product yields a compound that is more toxic compared to the parent compound. Furthermore, the reactions of Phase 1 enzymes generate very reactive damaging free radicals. Therefore it is necessary that this compound is further metabolized by the Phase 2 enzymes to avoid damage of the organism's proteins and genes. Phase 2 reactions consist of conjugating enzymes that normally adds a water-soluble molecule to the active site of the Phase 1 metabolite to facilitate its removal in urine or bile. These conjugates are usually less toxic and more water-soluble than their precursors. However, not all toxins undergo detoxification using the Phase 1 to Phase 2 route. Occasionally the parent toxins undergo detoxification directly through the Phase 2 route bypassing the Phase 1 route. Of all the Phase 2 enzymes, glutathione S-transferases (GSTs E.C. 2.5.1.18), the most abundant protein in the cytosolic fraction of the liver (Booth *et al.*, 1961; Wilce and Parker, 1994), is the most widely studied and major Phase 2 detoxification enzyme catalyzing the conjugation of cellular nucleophile glutathione with a wide range of endogenous or xenobiotic hydrophobic molecules (Armstrong, 1997; Hayes and Pulford, 1995; Mannervik and Danielson, 1988).

### **Glutathione**

Reduced glutathione (GSH), a tripeptide consisting of  $\gamma$ -glutamyl-cysteinyl-glycine can be conjugated in the initial step of mercapturic acid synthesis (George, 1994). Its cysteinyl residue provides a nucleophilic thiol important for the detoxication of electrophilic metabolites. Its net negative charge and overall hydrophilicity increases the aqueous solubility of the lipophilic